




Mark Malchoff, Lake Champlain Sea Grant Plattsburgh State Univ., Plattsburgh, NY



#### MRFSS B2 TRENDS



Personal communication from the NMFS, Fisheries Statistics and Economics Division, Silver Spring, MD




# **Hooking Mortality**

- **Wounding**
- Stress
- Depressurization or barotrauma



# Causes of Mortality: Hook Wounding

#### Wounding Mortality




Lowest wounding mortalities of sport-caught fish occur when related to hooking in the snout, maxillary, corner of mouth and cheek.

# Causes of Mortality: Angling Stress

#### **General Pattern**

- -muscular exertion
- -production of lactic acid (C<sub>3</sub>H<sub>6</sub>O<sub>3</sub>) in muscle tissues
- -increase in blood lactate
- -decrease in blood pH (blood becomes more acid- "blood acidosis")
- -fish has problems regulating exchange of ions (H+, CO<sub>3</sub>, etc.) between body tissues and the water

#### **Blood Lactate in rainbow trout**




Blood lactate concentration ([La<sup>-</sup>1)] in rainbow trout at rest (R) and after 0, 0.5, 1, and 4 hrs under control conditions (■), following exhaustive exercise (●) or exh. exercise plus 60 s of air exposure (○). (*From:* Ferguson and Tufts, 1992).

## Causes of Mortality Barotrauma

- Swimbladder overinflation/rupture
- Displacement of stomach or intestines
- Bulging eyes

More info at: http://edis.ifas.ufl.edu/SG047



## Research Issue Can hooking mortality be reduced?

Much of recent research has dealt with issues of


- artificial lures vs. bait
- hook style/size
- role of water temperature
- other variables

# **Hook Styles and Sizes**

Sproat
Size 2/0



Wide Gap Size 2/0



Circle Size 4/0



### Methods

(used by Malchoff and others)

- Series of angling trials (10 to 25 fish/ trial)
- All fish tagged and caged near angling site. Recorded hook style/size, hook removal status (left or removed) wound location, presence of bleeding, salinity and water temp.
- All fish recovered after 72 hours. Mortalities expressed as percentage of total number of fish in each trial.
- Bootstrapping to develop mean mortality w/ 95% C.I.
- Modeled mortality using logistic regression

## Field methods






## Thousands of fish released – how many live?

	Temp (oC)	Salinity (ppt)	Number of fish	Percent Mortality
Trial 1	27	29	26	0.0
Trial 2	23	23	31	6.5
Trial 3	23	25	26	3.8
Trial 4	22	32	7	0.0
Total			90	Tire State S
Mean mortality (%)				2.6
95% Confidence	Interval			0.6-7.0

## Hook Type vs. Wound Location

(summer flounder)



# **Logistic Regression Results**

Parameter	Probability
Leader status	<0.0001
Hook type	>0.05
Wound location	<0.0001
Bleeding	<0.0001
<b>Temperature in degrees Celsius</b>	<0.05
Length (mm)	>0.82
Bleeding*Hook Type (interaction)	<0.0001
Bleeding*Length (interaction)	<0.01

From Malchoff et al., 2002

# **Overall Summer Flounder Mortality**

	NY	VA – NC combined	Overall (total)
Mortality	14.6	6.1	9.5
95% CI	7.9 – 21.2	3.2 – 9.5	7.4 – 11.9
Sample	247	376	<b>623</b>

From Malchoff et al., 2002

### Other studies

- Circle hooks significant predictor of wound location in tuna. Bluefin c&r mortality estimated at 4% with circle hooks, and 28% on straight hooks (Skomal, Chase and Prince, 2002)
- Circle hooks resulted in lower mortality in 16 species in pelagic longline fishery (Falterman and Graves, 2002)
- "Careful release" (rolling hook out, cutting gangion) reduced discard mortality to 11% in Pacific halibut fishery (Trumble, Kaimmer, and Williams, 2002)
- Fewer striped bass were deeply hooked (10.6% vs. 45.6%) with circle hooks as compared to standard hooks in Chesapeake Bay (Lukacovic and Uphoff, 2002)
- Little difference treble hooks and single hooks on mortality in spotted seatrout (Duffy, 2002)
- Mortality in tautog was significantly higher in deep-water (11-17 m) vs. shallow water (≤10m) (Lucy and Arendt, 2002)

#### Recommendations

- Minimize stress
- Try circle hooks or barbless hooks to minimize wounding
- Set hook quickly if fishing live bait
- Use dehooker or other techniques to minimize handling
- Contact Sea Grant for additional advice

